arXiv:math/0201096v1 [math.DS] 11 Jan 2002

Elliptic Islands on Strictly Convex Billiards

Mário Jorge DIAS CARNEIRO, Sylvie OLIFFSON KAMPHORST
and Sônia PINTO DE CARVALHO
Departamento de Matemática, ICEx, UFMG 30.123–970, Belo Horizonte, Brasil. *

January 2002

Abstract

This paper addresses the question of genericity of existence of elliptic islands for the billiard map associated to strictly convex closed curves. More precisely, we study 2-periodic orbits of billiards associated to C^5 closed and strictly convex curves and show that the existence of elliptic islands is a dense property on the subset of those billiards having an elliptic 2-periodic point.

Our main tools are normal perturbations, the Birkhoff Normal Form for elliptic fixed points and Moser's Twist Theorem.

1 Introduction

Let α be a plane, closed, regular, strictly convex, oriented counterclockwise. Any curve with these properties may be parametrized by $\varphi \in [0, 2\pi]$, the angle between the tangent vector and a given direction The billiard problem on α consists in the free motion of a point particle in the plane region enclosed by α, being reflected elastically at the impacts with the boundary. Since the particle moves with constant velocity inside the region, the motion is completely determined by the point of reflection at α and the direction of movement immediately after each reflection. So, the angle φ, which locates the point of reflection, and the angle θ between the direction of motion and the normal to the boundary at the reflection point, may be used to describe the system.

The billiard model defines a map T from the annulus $\mathcal{A} = [0, 2\pi) \times (-\pi/2, \pi/2)$ into itself. If α is C^k, T is a C^{k-1}-diffeomorphism [7] preserving the measure $d\mu = R(\varphi) \cos\theta \, d\theta d\varphi$ [1], where $R(\varphi)$ is the radius of curvature of α at φ. $(\mathcal{A}, \mu, T)$ defines a two dimensional discrete dynamical system, whose orbits are given by $\{(\varphi_n, \theta_n) = T^n(\varphi_0, \theta_0), n \in Z\}$.

A point $(\varphi_0, \theta_0) \in \mathcal{A}$ is n-periodic if n is the smallest positive integer such that $T^n(\varphi_0, \theta_0) = (\varphi_0, \theta_0)$. It is called elliptic if the eigenvalues of $DT^n_{(\varphi_0,\theta_0)}$ are complex, with non vanishing real part. An elliptic island is a T^n-invariant subset, homeomorphic to a disk, surrounding a n-periodic elliptic point.

We study 2-periodic orbits of billiards associated to C^5 closed and strictly convex curves and show that the existence of elliptic islands is a dense property on the subset of those billiards having an elliptic 2-periodic point.

---

*email: carneiro@mat.ufmg.br, syok@mat.ufmg.br, sonia@mat.ufmg.br

It is well known that for strictly convex curves, at least C^6, the billiard map has invariant curves near the boundary of the phase space [2], and then it is not ergodic. Here we study the dynamical properties of the region of the phase space between these curves, when they exist, or the whole phase space, otherwise. The existence of islands assures the non-ergodicity of this region.

The layout of the paper is as follows: we will deal with plane, closed, regular, strictly convex, oriented counterclockwise curves. In sections 2 and 3 we analyze the geometrical and dynamical properties of the 2-periodic orbits. In section 4, we define normal perturbations and prove that the existence of elliptic or hyperbolic 2-periodic orbits is an open property. Sections 5 and 6 deal with the resonances and the existence of elliptic islands. In sections 7 we present a more topological description of the results obtained in the previous sections. We conclude this work with some comments and open questions. Our main tools are normal perturbations, the Birkhoff Normal Form for elliptic fixed points and Moser's Twist Theorem.

2 The Diameters of a Convex Curve

Let a curve α be parametrized by $\varphi \in [0, 2\pi]$, the angle between the tangent vector and a given direction, hereafter denoted by $\mathbf{x}$. Let $\tau(\varphi)$ and $\eta(\varphi)$ be the unitary tangent and (inward) normal vectors. Then $\alpha'(\varphi) = R(\varphi)\tau(\varphi)$. Clearly, $\tau(\varphi + \pi) = -\tau(\varphi)$ and $\eta(\varphi + \pi) = -\eta(\varphi)$.

It is known that any closed convex curve has at least a width and a diameter, characterized by the global maximum and minimum of the distance between points with parallel tangents (or with double normal) (see, for instance, [3]). We broaden this concept defining:

Definition 2.1 *φ_0 is a diameter of α if $\alpha(\varphi_0) - \alpha(\varphi_0 + \pi)$ is parallel to $\eta(\varphi_0)$.*

Proposition 2.2 *Let $l(\varphi) = |\alpha(\varphi) - \alpha(\varphi + \pi)|$. Then φ_0 is a diameter of α if and only if $\frac{dl}{d\varphi}(\varphi_0) = 0$*

Proof:

$$\begin{aligned} 2l\frac{dl}{d\varphi} &= 2\left\langle \alpha'(\varphi) - \alpha'(\varphi + \pi), \alpha(\varphi) - \alpha(\varphi + \pi) \right\rangle = \\ &= 2\left\langle R(\varphi)\tau(\varphi) - R(\varphi + \pi)\tau(\varphi + \pi), \alpha(\varphi) - \alpha(\varphi + \pi) \right\rangle = \\ &= 2\left\langle (R(\varphi) + R(\varphi + \pi))\tau(\varphi), \alpha(\varphi) - \alpha(\varphi + \pi) \right\rangle \end{aligned}$$

So

$$\frac{dl}{d\varphi}(\varphi_0) = 0 \Longleftrightarrow \left\langle \alpha(\varphi_0) - \alpha(\varphi_0 + \pi), \tau(\varphi_0) \right\rangle = 0$$

◊

Let $L_0 = l(\varphi_0)$, $R_0 = R(\varphi_0)$ and $R_\pi = R(\varphi_0 + \pi)$.

Proposition 2.3 *Let φ_0 be a diameter. If $L_0 - (R_0 + R_\pi) \neq 0$ then φ_0 is an isolated diameter.*

Proof:

$$\begin{aligned}\frac{d}{d\varphi}\left(2l\frac{dl}{d\varphi}\right) &= 2\left(\frac{dl}{d\varphi}\right)^2 + 2l\frac{d^2l}{d\varphi^2} = \\ &= 2\left(\langle (R'(\varphi)+R'(\varphi+\pi))\tau(\varphi) + (R(\varphi)+R(\varphi+\pi))\eta(\varphi), \alpha(\varphi)-\alpha(\varphi+\pi)\rangle + \right. \\ &\quad \left. + \langle (R(\varphi)+R(\varphi+\pi))\tau(\varphi), (R(\varphi)+R(\varphi+\pi))\tau(\varphi)\rangle\right)\end{aligned}$$

At the diameter φ_0:

$$2l(\varphi_0)\frac{d^2l}{d\varphi^2}(\varphi_0) = 2(-L_0(R_0+R_\pi) + (R_0+R_\pi)^2) = -2(R_0+R_\pi)(L_0-(R_0+R_\pi)).$$

and we have that if $L_0-(R_0+R_\pi)\neq 0$ then φ_0 is an isolated singular point of l. $\diamondsuit$

3 Dynamical Characterization of the Diameters of a Convex Curve

Let α be a C^k curve, $k\geq 2$, and T its associated billiard map.

If φ_0 is a diameter for α then $(\varphi_0, 0)$ is a 2-periodic orbit for T, and, as we have pointed out, any billiard on a strictly convex curve has at least two of those orbits. Its stability is given by the eigenvalues of:

$$DT^2_{(\varphi_0,0)} = \frac{1}{R_0R_\pi}\begin{pmatrix} L_0-R_\pi & L_0 \\ L_0-(R_0+R_\pi) & L_0-R_0\end{pmatrix}\begin{pmatrix} L_0-R_0 & L_0 \\ L_0-(R_0+R_\pi) & L_0-R_\pi\end{pmatrix}$$

it follows that:

1. $(\varphi_0, 0)$ is hyperbolic if
   - $L_0-(R_0+R_\pi)>0$ or
   - $L_0-(R_0+R_\pi)<0$ and $(L_0-R_0)(L_0-R_\pi)<0$.
2. $(\varphi_0, 0)$ is elliptic if $L_0-(R_0+R_\pi)<0$ and $(L_0-R_0)(L_0-R_\pi)>0$.
3. $(\varphi_0, 0)$ is parabolic if
   - $L_0-(R_0+R_\pi)=0$ or
   - $L_0-(R_0+R_\pi)<0$ and $(L_0-R_0)(L_0-R_\pi)=0$.

We will say that a diameter is hyperbolic, elliptic or parabolic if the associated 2-periodic orbit is, respectively, hyperbolic, elliptic or parabolic.

A nice consequence [4] of this characterization is that, for $k\geq 3$, the largest diameter of a strictly convex curve, if isolated, is hyperbolic, due to the fact that $\frac{d^2l}{d\varphi^2}(\varphi_0) = -\frac{(R_0+R_\pi)(L_0-(R_0+R_\pi))}{L_0} < 0$.

The smallest one, however, can be elliptic or hyperbolic. If $R_0=R_\pi$ (which happens, for instance, for the ellipse), it is elliptic. Nevertheless, as we show in the last section, there are strictly convex curves such that all the diameters are hyperbolic.

When φ_0 is elliptic, the eigenvalues of $DT^2_{(\varphi_0,0)}$ are given by $e^{\pm i\gamma} = \frac{\mathrm{tr}(DT^2)}{2} \pm i\sqrt{1-(\frac{\mathrm{tr}(DT^2)}{2})^2}$ and so

$$\cos\gamma = 2\frac{(L_0-R_0)(L_0-R_\pi)}{R_0R_\pi} - 1\ . \tag{1}$$



Figure 1: Definition of $h(\varphi)$

4 Normal Perturbations of a Convex Curve

Let α C^k, with $k \geq 4$, when parametrized by $\varphi \in [0, 2\pi]$, the angle between the tangent vector and $\mathbf{x}$. Using this direction and its perpendicular as a reference frame, the unitary tangent and (inward) normal vectors are written, respectively, as $\tau(\varphi) = (\cos\varphi, \sin\varphi)$ and $\eta(\varphi) = (-\sin\varphi, \cos\varphi)$. So, the normal bundle $(\alpha(\varphi), \eta(\varphi))$ is also C^k and $\eta(\varphi)$ is C^∞.

A normal perturbation of α is a curve β given by $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$, where λ is a C^k, 2π-periodic function. Observe that in general φ does not represent the angle between the tangent vector of β and the fixed direction $\mathbf{x}$.

We will use the norm $|\lambda|_2 = \max\limits_{\varphi \in [0,2\pi]} \{|\lambda(\varphi)|, |\lambda'(\varphi)|, |\lambda''(\varphi)|\}$.

Proposition 4.1 *If $|\lambda|_2$ is small enough then β is C^k, closed, regular and strictly convex.*

Proof: : β is closed because λ is 2π-periodic. α is C^k, λ is C^k and η is C^∞. So, β is C^k. $\beta' = \alpha' + \lambda'\eta - \lambda\tau$. So, if α is regular, for ϵ small enough, β is regular. $\beta'' \times \beta' = (\alpha'' \times \alpha') + \lambda'(\alpha'' \times \eta) - \lambda(\alpha'' \times \tau) + (\lambda'' - \lambda)(\eta \times \alpha') + (\lambda(\lambda'' - \lambda) + 2\lambda')(\tau \times \eta)$. So, if α is strictly convex, for ϵ small enough, β is strictly convex. ♢

Let us call $\vec{t}(\varphi)$ and $\vec{n}(\varphi)$ the unitary tangent and (inward) normal vectors and $R_\alpha(\varphi)$ and $R_\beta(\varphi)$ the radius of curvature of α and β at φ, respectively. Then:

$$\begin{aligned}
\vec{t} &= \frac{(R_\alpha - \lambda)\tau + \lambda'\eta}{((R_\alpha - \lambda)^2 + (\lambda')^2)^{1/2}} \\
\vec{n} &= \frac{-\lambda'\tau + (R_\alpha - \lambda)\eta}{((R_\alpha - \lambda)^2 + (\lambda')^2)^{1/2}} \\
R_\beta &= \frac{((R_\alpha - \lambda)^2 + (\lambda')^2)^{3/2}}{(R_\alpha - \lambda)(R_\alpha - \lambda + \lambda'') - \lambda'(R'_\alpha - 2\lambda')} \qquad (2)
\end{aligned}$$

Let $h(\varphi)$ be the angle between $\vec{t}(\varphi)$ and the given direction that defines φ, as in fig. 1. It is easy to see that

$$h(\varphi) = \varphi + \arctan \frac{\lambda'}{R_\alpha - \lambda}.$$

According to proposition 2.2, the diameters of β are such that the derivative of

$$l_\beta(\varphi) = |\beta(\varphi) - \beta(h^{-1}(h(\varphi) + \pi))|$$

vanishes.

Lemma 4.2 *If $|\lambda|_2$ is sufficiently small then h is a C^2-diffeomorphism.*

Proof: : As α is closed, $0 < c_1 \leq R_\alpha$. Then, for $|\lambda|_2$ sufficiently small, $R_\alpha - \lambda > 0$ and h is well defined. α and β being at least C^4 then λ is at least C^4, and so h is at least C^3.

Let s be the arclength parameter for β. Then

$$\frac{dh}{d\varphi} = \frac{dh}{ds}\frac{ds}{d\varphi} = k_\beta|\frac{d\beta}{d\varphi}| \geq c_2 > 0.$$

Then h is C^1 and invertible, and h^{-1} has continuous derivative.

$$\frac{d^2h}{d\varphi^2} = \frac{dk_\beta}{ds}|\frac{d\beta}{d\varphi}|^2 + k_\beta \frac{< \frac{d^2\beta}{d\varphi^2}, \frac{d\beta}{d\varphi} >}{|\frac{d\beta}{d\varphi}|}$$

so h is C^2. Taking $y = h(\varphi)$ we have

$$\frac{d^2h^{-1}}{dy^2}(y) = -\frac{\frac{d^2h}{d\varphi^2}(\varphi)}{(\frac{dh}{d\varphi}(\varphi))^3}$$

which shows that h is a C^2-diffeomorphism. ♢

Lemma 4.3 *With the same hypothesis of lemma 4.2, $f(\varphi) = h^{-1}(h(\varphi) + \pi)$ and $I_\pi(\varphi) = \varphi + \pi$ are C^2 close.*

Proof: : Let $|\lambda|_2 < \epsilon$ and $\frac{1}{c_1} < |R_\alpha - \lambda| < c_2$, then $|h(\varphi) - \varphi| = |\arctan \frac{\lambda'}{R_\alpha - \lambda}| \leq |\frac{\lambda'}{R_\alpha - \lambda}| < c_1\epsilon$

Let $c_3 = \max|\frac{dh^{-1}}{dy}|$.

$$|f(\varphi) - I_\pi(\varphi)| = |h^{-1}(h(\varphi) + \pi) - h^{-1}(h(\varphi + \pi))| \leq c_3|h(\varphi) + \pi - h(\varphi + \pi)| =$$

$$= c_3|h(\varphi) - \varphi + \varphi + \pi - h(\varphi + \pi)| \leq c_3(|h(\varphi) - \varphi| + |\varphi + \pi - h(\varphi + \pi)|) < 2c_1c_3\epsilon$$

Let $0 < \frac{1}{c_4} < |R'_\alpha - \lambda'| < c_5$ and $\Delta(\varphi) = \arctan \frac{\lambda'}{R_\alpha - \lambda}$. Now, $\frac{dh}{d\varphi} = 1 + \frac{d\Delta}{d\varphi}$ and

$$|\frac{d\Delta}{d\varphi}| \leq |\frac{d}{d\varphi}\frac{\lambda'}{R_\alpha - \lambda}| = |\frac{\lambda''(R_\alpha - \lambda) - \lambda'(R'_\alpha - \lambda')}{(R_\alpha - \lambda)^2}| < c_1^2c_2c_5\epsilon \tag{3}$$

Taking $y = h(\varphi)$,

$$\begin{aligned}
\left|\frac{df}{d\varphi} - 1\right| &= \left|\frac{dh^{-1}}{dy}(h(\varphi) + \pi)\frac{dh}{d\varphi}(\varphi) - 1\right| = \left|\frac{1 + \frac{d\Delta}{d\varphi}(\varphi)}{1 + \frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi))} - 1\right| = \\
&= \left|\frac{1 + \frac{d\Delta}{d\varphi}(\varphi) + \frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi)) - \frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi))}{1 + \frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi))} - 1\right| = \\
&= \left|\frac{\frac{d\Delta}{d\varphi}(\varphi) - \frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi))}{1 + \frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi))}\right| \leq \\
&\leq \left|\frac{dh^{-1}}{d\varphi}(h(\varphi) + \pi))\right| \left|\frac{d\Delta}{d\varphi}(\varphi)\right| + \left|\frac{d\Delta}{d\varphi}(h^{-1}(h(\varphi) + \pi))\right| < 2c_3c_1^2c_2c_5\epsilon
\end{aligned}$$

Finally,

$$|\frac{d^2f}{d\varphi^2}(\varphi) - 0| = |\frac{d}{d\varphi}(\frac{dh^{-1}}{dy}(h(\varphi)+\pi)\frac{dh}{d\varphi}(\varphi))| \leq$$
$$|\frac{d^2h^{-1}}{dy^2}(h(\varphi)+\pi)||\frac{dh}{d\varphi}|^2 + |\frac{dh^{-1}}{dy}(h(\varphi)+\pi)||\frac{d^2h}{d\varphi^2}(\varphi))|$$

$|\frac{dh}{d\varphi}|$ and $|\frac{dh^{-1}}{dy}(h(\varphi)+\pi)|$ are bounded from above and bellow because h is a C^2 diffeomorphism on $[0, 2\pi]$.

$$|\tfrac{d^2h}{d\varphi^2}(\varphi))| = |\tfrac{d^2\Delta}{d\varphi^2}(\varphi))| \leq$$
$$|\tfrac{\lambda'''(R_\alpha-\lambda)-\lambda'(R''_\alpha-\lambda'')}{(R_\alpha-\lambda)^2+(\lambda')^2}| + 2|\tfrac{(\lambda''(R_\alpha-\lambda)-\lambda'(R'_\alpha-\lambda'))(R'_\alpha-\lambda')}{(R_\alpha-\lambda)((R_\alpha-\lambda)^2+(\lambda')^2)}| + 2|\tfrac{\lambda'}{R_\alpha-\lambda}||\tfrac{d\Delta}{d\varphi}(\varphi)|$$

For ϵ small enough, $0 < \frac{1}{c_1} < |R_\alpha - \lambda| < c_2$, $|R'_\alpha - \lambda'| < c_5$, $|R''_\alpha - \lambda''| < c_6$ and $(R_\alpha - \lambda)^2 + (\lambda')^2 = |\beta'(\varphi)|^2 \geq c_7 > 0$, since β is regular. Together with equation 3, we have that $|\frac{d^2h}{d\varphi^2}(\varphi))| < K\epsilon$, for a constant K.

$$|\frac{d^2h^{-1}}{dy^2}(h(\varphi)+\pi)| = |\frac{\frac{d^2h}{d\varphi^2}(h^{-1}(h(\varphi)+\pi))}{(\frac{dh}{d\varphi}(h^{-1}(h(\varphi)+\pi)))^3}| < K\epsilon \tag{4}$$

for a constant K. So, $|\frac{d^2f}{d\varphi^2}(\varphi)| < C\epsilon$. ♢

Lemma 4.4 *Let* $l_\alpha(\varphi) = |\alpha(\varphi) - \alpha(\varphi+\pi)|$ *and* $l_\beta(\varphi) = |\beta(\varphi) - \beta(h^{-1}(h(\varphi)+\pi))|$. *If* $|\lambda|_2$ *is small enough then* l_α *and* l_β *are* C^2 *close.*

Proof: : If $|\lambda|_2$ is sufficiently small then α and β are C^2 close. So $\alpha \circ I_\pi$ and $\beta \circ f$ are C^2 close and the result follows. ♢

Proposition 4.5 *Suppose that* α *has an isolated diameter at* φ_0, *with* $\frac{dl_\alpha}{d\varphi}(\varphi_0) = 0$ *and* $\frac{d^2l_\alpha}{d\varphi^2}(\varphi_0) > 0$ *(or* < 0*). Let* δ *be such that* $\frac{d^2l_\alpha}{d\varphi^2}(\varphi) > 0$ *(or* < 0*) for* $\varphi \in [\varphi_0 - \delta, \varphi_0 + \delta]$. *Then there exists* ϵ_0 *such that if* $|\lambda|_2 < \epsilon_0$ *then any normal perturbation* $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$ *has an isolated diameter* $\varphi_1 \in (\varphi_0 - \delta, \varphi_0 + \delta)$, *with* $\frac{d^2l_\beta}{d\varphi^2}(\varphi_1) > 0$ *(or* < 0*). Furthermore,* $\epsilon_0 \to 0$ *if* $\delta \to 0$.

Proof: : The proof follows immediately from the results above and from the transversality of the zero of $\frac{dl_\alpha}{d\varphi}$.
♢

Observe that rotating the axis, if necessary, we can always suppose that the diameter of α is at $\varphi = 0$.

Let, as above, $L_0 = l_\alpha(0)$, $R_0 = R_\alpha(0)$, and $R_\pi = R_\alpha(\pi)$. An obvious consequence is:

Corollary 4.6 *If* 0 *is a hyperbolic diameter of* α, *with* $L_0 - (R_0 + R_\pi) > 0$ *(meaning that* $\frac{d^2l_\alpha}{d\varphi^2}(0) < 0$*) then every normal perturbation* β *with* $|\lambda|_2 < \epsilon_0$ *has a hyperbolic diameter close to* 0.

Proposition 4.7 *Let* ϵ_0 *and* δ *be as in Proposition 4.5 and suppose* 0 *is an elliptic (resp. hyperbolic) diameter for* α, *i.e.* $L_0 - (R_0 + R_\pi) < 0$ *and* $(L_0 - R_0)(L_0 - R_\pi) > 0$ *(resp.* $(L_0 - R_0)(L_0 - R_\pi) < 0$*). There exists* $\epsilon_1 \leq \epsilon_0$ *such that if* $|\lambda|_2 < \epsilon_1$ *then any normal perturbation* $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$ *has an elliptic (resp. hyperbolic) diameter at* $\varphi_1 \in (\varphi_0 - \delta, \varphi_0 + \delta)$.

Proof: If $|\lambda|_2 < \epsilon_0$ then, by Proposition 4.5, β has an isolated diameter at $\varphi_1 \in (\varphi_0 - \delta, \varphi_0 + \delta)$ with $l_\beta(\varphi_1) - (R_\beta(\varphi_1) + R_\beta(h^{-1}(h(\varphi_1) + \pi)) < 0$. Since α and β are C^2-close, R_α and R_β are C^0-close. This, together with lemma 4.3 and $(L_0 - R_0)(L_0 - R_\pi) > 0$, gives that there exists $\epsilon_1 \le \epsilon_0$ such that if $|\lambda|_2 < \epsilon_1$ then $(l_\beta(\varphi_1) - R_\beta(\varphi_1))(l_\beta(\varphi_1) - R_\beta(h^{-1}(h(\varphi_1) + \pi))) > 0$. ♢

The hyperbolic case $(L_0 - R_0)(L_0 - R_\pi) < 0$ is analogous.

Remark: Propositions 4.5 and 4.7 and corollary 4.6 only express, in the context of billiard maps, the well known fact of openness of existence of elliptic or hyperbolic fixed points.

5 Resonances

We will assume that the curve α has an elliptic diameter with eigenvalues $e^{\pm i\gamma}$. Without any loss of generality we can assume that this diameter is located at $\varphi = 0$.

Definition 5.1 *An elliptic diameter is resonant if its eigenvalues satisfy* $(e^{\pm i\gamma})^j = 1$, *for* $j = 2, 3, 4$, *i.e., if* $\gamma = \frac{1}{2}\pi$, $\frac{2}{3}\pi$, $\frac{4}{3}\pi$, $\frac{3}{2}\pi$. *Otherwise, it is non-resonant.*

Proposition 5.2 *Given a C^k strictly convex curve α, $k \ge 4$, with a resonant elliptic diameter $\varphi = 0$, there is a C^k strictly convex curve β, C^2-close to α, such that β has a non-resonant elliptic diameter.*

Proof: For ϵ_1 as in Proposition 4.7 consider the normal perturbations $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$, with $|\lambda|_2 < \epsilon_1$ and $\lambda'(0) = \lambda'(\pi) = 0$, which implies that $\varphi = 0$ is also an elliptic diameter for β.

Let $\lambda(0) = \lambda_0$, $\lambda(\pi) = \lambda_\pi$, $\lambda''(0) = \lambda_0''$, $\lambda''(\pi) = \lambda_\pi''$. We have that $l_0 = l_\beta(0) = |\beta(0) - \beta(\pi)| = L_0 - \lambda_0 - \lambda_\pi$ and, by equation (2),

$$r_0 = R_\beta(0) = \frac{(R_0 - \lambda_0)^2}{R_0 - \lambda_0 + \lambda_0''} = R_0 - \lambda_0 - \lambda_0'' \frac{R_0 - \lambda_0}{R_0 - \lambda_0 + \lambda_0''}$$

and

$$r_\pi = R_\beta(\pi) = \frac{(R_\pi - \lambda_\pi)^2}{R_\pi - \lambda_\pi + \lambda_\pi''} = R_\pi - \lambda_\pi - \lambda_\pi'' \frac{R_\pi - \lambda_\pi}{R_\pi - \lambda_\pi + \lambda_\pi''}$$

So, for $|\lambda|_2$ small enough, these normal perturbations have an elliptic diameter with eigenvalues $e^{\pm i\gamma(\beta)}$ satisfying, by equation (1)

$$\begin{aligned} \cos\gamma(\beta) &= 2\,\frac{(l_0 - r_0)(l_0 - r_\pi)}{r_0 r_\pi} - 1 = \\ &= 2\,\frac{(L_0 - R_0 - \lambda_\pi + \lambda_0''\Delta_0)(L_0 - R_\pi - \lambda_0 + \lambda_\pi''\Delta_\pi)}{(R_0 - \lambda_0 - \lambda_0''\Delta_0)(R_\pi - \lambda_\pi - \lambda_\pi''\Delta_\pi)} - 1 = \\ &= F(\lambda_0, \lambda_\pi, \lambda_0', \lambda_\pi') \end{aligned}$$

where

$$\Delta_a = \frac{R_a - \lambda_a}{R_a - \lambda_a + \lambda_a''} \ .$$

Clearly $F(0,0,0,0) = 2\frac{(L_0 - R_0)(L_0 - R_\pi)}{R_0 R_\pi} - 1$ gives the eigenvalues of the elliptic diameter of α.

Moreover

$$\begin{aligned}
\frac{\partial F}{\partial \lambda_0}(0,0,0,0) &= 2\frac{(L_0-R_0)(L_0-R_0-R_\pi)}{R_0^2 R_\pi} \neq 0\\
\frac{\partial F}{\partial \lambda_\pi}(0,0,0,0) &= 2\frac{(L_0-R_\pi)(L_0-R_0-R_\pi)}{R_\pi^2 R_0} \neq 0\\
\frac{\partial F}{\partial \lambda_0''}(0,0,0,0) &= 2\frac{(L_0-R_\pi)L_0}{R_0^2 R_\pi} \neq 0\\
\frac{\partial F}{\partial \lambda_\pi''}(0,0,0,0) &= 2\frac{(L_0-R_0)L_0}{R_\pi^2 R_0} \neq 0
\end{aligned}$$

which shows that F is a non constant rational function and the resonance conditions of definition 5.1 are fulfilled only on a union of codimension 1 submanifolds. Hence it is enough to choose a normal perturbation with λ small, $\lambda'(0) = \lambda'(\pi) = 0$ and outside this set to obtain a curve β, close to α, with a non-resonant elliptic diameter. ♢

Proposition 5.3 *If 0 is a non-resonant elliptic diameter for a strictly convex C^k curve α, $k \geq 3$, then there exists ϵ_2 such that every normal perturbation β, with $|\lambda|_2 < \epsilon_2$, has a non-resonant elliptic diameter.*

Proof: : As 0 is a non-resonant elliptic diameter for α then

$$\begin{aligned}
\cos\gamma(\alpha) &= 2\,\frac{(l_\alpha(0)-R_\alpha(0)\,(l_\alpha(0)-R_\alpha(\pi))}{R_\alpha(0)R_\alpha(\pi)} - 1 = \\
&= 2\,\frac{(L_0-R_0)(L_0-R_\pi)}{R_0R_\pi} - 1 \neq 0, 1, -1, \cos\frac{2}{3}\pi, \cos\frac{4}{3}\pi
\end{aligned}$$

For ϵ_1 as in Proposition 4.7, let $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$, with $|\lambda|_2 < \epsilon_1$. Then β has an elliptic diameter at $\varphi_1 \in (-\delta, \delta)$ with eigenvalues determined by

$$\cos\gamma(\beta) = 2\,\frac{(l_\beta(\varphi_1)-R_\beta(\varphi_1))(l_\beta(\varphi_1)-R_\beta(h^{-1}(h(\varphi_1)+\pi))}{R_\beta(\varphi_1)R_\beta(h^{-1}(h(\varphi_1)+\pi))} - 1.$$

Since α and β are C^k, $k \geq 3$, then R_α and R_β are C^1 close. This, together with lemmas 4.3 and 4.4, gives the desired result. ♢

6 Elliptic Islands

In order to seek after islands, let us change coordinates from (φ, θ) to (s, p), where s is the arclenght parameter for the curve α and $p = \sin\theta$. The advantage of those new coordinates is that the billiard map T preserves the area $ds \wedge dp$.

Let us now suppose that, for $k \geq 5$, the C^k curve α has a non resonant elliptic diameter at $s = 0$, so that $(0, 0)$ is a non resonant elliptic fixed point of T^2.

By means of a linear area preserving coordinate change, and complexification, $T^2(s, p)$ can be brought into the form $e^{i\gamma}z + g(z, \overline{z})$ where $z, \overline{z}$ are complex and g vanishes with its first derivative at $z = 0$. This map has then the Birkhoff Normal Form $e^{i(\gamma+\tau_1(\alpha)|\zeta|^2)}\zeta + h(\zeta, \overline{\zeta})$, where $h(\zeta, \overline{\zeta}) = \mathcal{O}(|\zeta|^4)$. Moreover, if the first

Birkhoff coefficient $\tau_1(\alpha) \neq 0$ then, by Moser's Twist Theorem [6], $\zeta = 0$ is a stable fixed point, i.e., $(0,0)$ has T^2-invariant curves surrounding it which means that there is an elliptic island of positive measure.

Hence, by assuming $\tau_1(\alpha) = 0$, the goal is show that small C^2-normal perturbations of α may have non-zero Birkhoff coefficients.

As Moeckel showed in [5], τ_1 can be calculated using complex area preserving coordinates diagonalizing the linear part of the map, such that the third jet of T^2 takes the form:

$$Z = e^{i\gamma}(z + c_{20}z^2 + c_{11}z\overline{z} + c_{02}\overline{z}^2 + c_{30}z^3 + c_{21}z^2\overline{z} + c_{12}z\overline{z}^2 + c_{03}\overline{z}^3)$$

then the Birkhoff coefficient is given by

$$\tau_1 = \frac{1}{i}\left(c_{21} + 2|c_{20}|^2\left(\frac{2e^{i\gamma}+1}{e^{i\gamma}-1} + \frac{1}{e^{3i\gamma}-1}\right)\right) .$$

The first step is, then, to determine the third jet of T^2 at $(0,0)$. Although we are not able to write T explicitly, we have that (see, for instance, [7]):

$$\begin{aligned}
\frac{\partial s_1}{\partial s_0} &= \frac{l(s_0,p_0) - R(s_0)\cos\theta(p_0)}{R(s_0)\cos\theta(p_1)}\\
\frac{\partial s_1}{\partial p_0} &= \frac{l(s_0,p_0)}{\cos\theta(p_0)\cos\theta(p_1)}\\
\frac{\partial p_1}{\partial s_0} &= \frac{l(s_0,p_0) - R(s_0)\cos\theta(p_0) - R(s_1)\cos\theta(p_1)}{R(s_0)R(s_1)}\\
\frac{\partial p_1}{\partial p_0} &= \frac{l(s_0,p_0) - R(s_1)\cos\theta(p_1)}{R(s_1)\cos\theta(p_0)}
\end{aligned}$$

where $(s_1, p_1) = T(s_0, p_0)$.

Using the Chain Rule and with the aid of the software Maple [1], we have calculated the coefficients of the third-jet $J_3\,T^2_{(0,0)}(s_0,p_0)$ and obtained the above coefficients c_{ij}. They depend on L_0, $R_0 = R(0)$, $R_\pi = R(s_1)$, $\frac{dR}{ds}(0)$, $\frac{dR}{ds}(s_1)$, $\frac{d^2R}{ds^2}(0)$, $\frac{d^2R}{ds^2}(s_1)$, where $(s_1, 0) = T(0,0)$. Moreover, c_{21} depends linearly on $\frac{d^2R}{ds^2}(0)$ and $\frac{d^2R}{ds^2}(s_1)$ and c_{20} does not depend on those second derivatives.

Using these calculations we can prove:

Proposition 6.1 *Let α be a C^k strictly convex curve, $k \geq 5$, with a non-resonant elliptic diameter at $s = 0$, and such that $\tau_1(\alpha) = 0$. Then there is a C^k strictly convex curve β, C^2-close to α, such that β has a non-resonant elliptic diameter, with $\tau_1(\beta) \neq 0$.*

Proof: : Let $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$ be a normal perturbation of α, with a third order contact with α at $\varphi(0) = 0$ and $\varphi(s_1) = \pi$. This means that $\lambda(0) = \lambda(\pi) = \lambda'(0) = \lambda'(\pi) = \lambda''(0) = \lambda''(\pi) = \lambda'''(0) = \lambda'''(\pi) = 0$ and then, β has a non-resonant elliptic diameter at $\varphi = 0$

Let σ be the arclenght parameter for β and $(\sigma_1, 0) = T^2_\beta(0,0)$. The third order contact implies that the elliptic diameters coincide, with the same length L_0 and

$$R_\beta(0) = R_\alpha(0) = R_0, \;\; R_\beta(\sigma_1) = R_\alpha(s_1) = R_\pi,$$

---

[1] the program may be found at www.mat.ufmg.br/∼syok

$$\frac{dR_\beta}{d\sigma}(0) = \frac{dR_\alpha}{ds}(0), \quad \frac{dR_\beta}{d\sigma}(\sigma_1) = \frac{dR_\alpha}{ds}(s_1),$$

$$\frac{dR_\beta^2}{d\sigma^2}(0) = \frac{dR_\alpha^2}{ds^2}(0) - \frac{d^4\lambda}{d\varphi^4}(0), \quad \frac{dR_\beta^2}{d\sigma^2}(\sigma_1) = \frac{dR_\alpha^2}{ds^2}(s_1) - \frac{d^4\lambda}{d\varphi^4}(\pi).$$

Then $\gamma(\beta) = \gamma(\alpha)$, $c_{20}(\beta) = c_{20}(\alpha)$ and

$$c_{21}(\beta) = -\frac{i}{16}\left[\frac{L_0(L_0 - R_\pi)}{(L_0 - R_0)(L_0 - R_0 - R_\pi)}\right]\frac{d^4\lambda}{d\varphi^4}(0) + ib(L_0, R_0, R_\pi)\frac{d^4\lambda}{d\varphi^4}(\pi) + c_{21}(\alpha),$$

where b is a C^{k-4} function.

Hence

$$\tau_1(\beta) = -\frac{1}{16}\left[\frac{L_0(L_0 - R_\pi)}{(L_0 - R_0)(L_0 - R_0 - R_\pi)}\right]\frac{d^4\lambda}{d\varphi^4}(0) + b(L_0, R_0, R_\pi)\frac{d^4\lambda}{d\varphi^4}(\pi) + \tau_1(\alpha).$$

As $\sigma = 0$ is a non-resonant elliptic diameter for α and β, then $L_0 \neq 0$, $L_0 - R_0 \neq 0$, $L_0 - R_\pi \neq 0$ and $L_0 - R_0 - R_\pi \neq 0$. Therefore, by choosing $\frac{d^4\lambda}{d\varphi^4}(0) \neq 0$ and $\frac{d^4\lambda}{d\varphi^4}(\pi) = 0$ we obtain a perturbation λ such that $\tau_1(\beta) \neq 0$.

7 Density of Elliptic Islands

Let α be a closed, regular, strictly convex plane curve and T_α its associated billiard map. Planar rigid motions do not change the geometrical features of α and then do not change the dynamical characteristics of T_α. This defines an equivalence relation on the set of curves.

Let $\mathcal{C}$ be the set of equivalent classes, denoted by $[\alpha]$, of closed, regular, strictly convex plane curves α, that are C^5 when parametrized by the angle φ.

Given a representative α of an equivalent class $[\alpha] \in \mathcal{C}$, the normal bundle $(\alpha(\varphi), \eta(\varphi))$ is then C^5. Given $\epsilon > 0$, let us consider a tubular neighbourhood $N_\epsilon(\alpha) = \{\alpha(\varphi) + \lambda\eta(\varphi), 0 \leq \varphi < 2\pi, -\epsilon < \lambda < \epsilon\} \simeq S^1 \times (-\epsilon, \epsilon)$ and the canonical projection $\Pi : S^1 \times (-\epsilon, \epsilon) \to S^1$ that projects $N_\epsilon(\alpha)$ onto the image of α.

Definition 7.1 *$[\beta] \in \mathcal{C}$ is ϵ-close to $[\alpha] \in \mathcal{C}$ if:*

1. *there exists $\beta \in [\beta]$ such that the image of β is in $N_\epsilon(\alpha)$, for $\alpha \in [\alpha]$.*
2. *the restriction of Π to the image of β is a diffeomorphism.*

As a consequence, β can be written as $\beta(\varphi) = \alpha(\varphi) + \lambda(\varphi)\eta(\varphi)$, with λ periodic.

Definition 7.2 *$[\beta] \in \mathcal{C}$ is ϵ-C^2-close to $[\alpha] \in \mathcal{C}$ if $[\beta]$ is ϵ-close to $[\alpha]$ and $|\lambda|_2 < \epsilon$.*

In this context, Propositions 4.7 and 5.3 can be rewritten as:

Proposition 7.3 *Let $\mathcal{E} \subset \mathcal{C}$ be the subset of equivalent classes of curves with an elliptic diameter. Then $\mathcal{E}$ is open in $\mathcal{C}$.*

Proposition 7.4 *Let $\mathcal{R} \subset \mathcal{E}$ be the subset of equivalent classes of curves with a non resonant diameter. Then $\mathcal{R}$ is open in $\mathcal{C}$.*

Propositions 5.2 and 6.1 leads to:

Theorem 1 *Any curve on an equivalent class of $\mathcal{E}$ can be approximated by curves such that the associated billiard map has an elliptic island of positive measure.*

Proof: : Let α be a representative of $[\alpha] \in \mathcal{E}$. So, α has an elliptic diameter at, we can suppose, $\varphi = 0$. There are three possible cases:

1. 0 is a non resonant diameter with $\tau_1(\alpha) \neq 0$.
2. 0 is a non resonant diameter with $\tau_1(\alpha) = 0$.
3. 0 is a resonant diameter.

In the first case, there is nothing to prove.

If 0 is a non resonant diameter with $\tau_1(\alpha) = 0$ then, by Proposition 6.1, α can be approximated by a sequence β_n such that for each n, β_n has a non resonant diameter with $\tau_1(\beta_n) \neq 0$.

If 0 is a resonant diameter, by Proposition 5.2, α can be approximated by a sequence β_n such that for each n, β_n has a non resonant diameter. The same reasoning used on the second case leads to the result. ♢

8 Conclusions and Comments

Although we can prove the density of elliptic islands surrounding orbits associated to diameters, many questions remain open.

The existence of islands is not equivalent to a non-vanishing first Birkhoff coefficient. For instance, any Birkhoff coefficient different from zero assures, for a sufficiently differentiable curve and avoiding higher order resonances, the existence of such islands. However one may ask if there is a strictly convex C^∞ curve with an elliptic diameter such that all the Birkhoff coefficients vanishes. If so, how this fact translates into the dynamics of the associated billiard map?

It may also happen that all the isolated diameters of a given curve are hyperbolic or parabolic. This is the case, for instance, of an ellipse with half-axes 1 and $\sqrt{2}$, which has only two diameters: the largest axis, which is hyperbolic, and the smallest one, which is parabolic. However, due to the integrability of the system, the parabolic orbit has an island. Kozlov, in [4] constructs a strictly convex curve, as differentiable as desired, that has exactly three diameters, all of them hyperbolic.

Even when all the diameters are hyperbolic, the examples suggest that a higher period trajectory is elliptic, if the curve is strictly convex. This is the case for the billiard associated to $x(t) = \cos t$, $y(t) = 3/(2 - \sin t)$. It has exactly two diameters, both hyperbolic. The numerical investigation of the phase-space of its associated



Figure 2: The curve $x(t) = \cos t$, $y(t) = 3/(2 - \sin t)$ with a period 3 trajectory and the phase space of the associated billiard map.

billiard map shows the existence of a pair of 3-periodic orbits, corresponding to a same triangular trajectory followed in both senses, and its islands (see figure 2). One may wonder if there is always an elliptic periodic orbit for any sufficiently differentiable and strictly convex curve.

If there are elliptic periodic orbits, can they have any period? In the example above only elliptic orbits of period 3 or multiples of 3, belonging to the same island, are visible.

Anyway, if there are elliptic orbits of higher period and they are non-resonant, we believe that our method of normal perturbations with a third order contact at the points of reflection of the periodic trajectory with the boundary will give the results we obtained for the 2-periodic orbit.

References

[1] G.D.Birkhoff: Dynamical Systems. Providence, RI: A. M. S. Colloquium Publications 1966 (Original ed. 1927)

[2] R. Douady: Applications du théorème des tores invariants. Thèse de 3ème Cycle, Univ. Paris VII (1982)

[3] A.Katok, B.Hasselblat: Introduction to the Modern Theory of Dynamical Systems. Univ. Cambridge Press 1997.

[4] V.V.Kozlov: Two-link billiard trajectories: extremal properties and stability, J.Appl.Maths Mechs, **64/6**, 903-907 (2000).

[5] R.Moeckel: Generic bifurcations of the twist coefficient, Erg.Th.Dyn.Syst. **10**, 185-195 (1990)

[6] J.Moser: *Stable and random motions in dynamical systems*, PUP, Princeton, 1973.

[7] J-M.Strelcyn: Plane Billiards as Smooth Dynamical Systems with Singularities. In: Katok, Strelcyn, in coll. with Ledrappier, Przytycki, LNM **1222**. Berlin, Heidelberg, New York: Springer-Verlag 1986